arXiv:math/9907191v1 [math.DG] 30 Jul 1999

A STEREOLOGICAL VERSION OF THE GAUSS-BONNET FORMULA

X. GUAL-ARNAU AND J.J. NUÑO-BALLESTEROS

1. Introduction

Let $D \subset S \subset \mathbb{R}^3$ be a domain with boundary in an orientable smooth surface S in $\mathbb{R}^3$. Then the Gauss-Bonnet formula gives a method to compute the Euler-Poincaré characteristic of D in terms of the Gauss curvature K of S, and the geodesic curvature k_g of ∂D in S:

$$2\pi\chi(D) = \int_D K + \int_{\partial D} k_g.$$

This can be found in any elementary book of Classical Differential Geometry. In this paper, we give a stereological version of this formula, in the following sense: let $u \in S^2$ and denote by $\pi_{u,\lambda}$ the plane orthogonal to u given by the equation $\langle x, u\rangle = \lambda$. When λ varies on $\mathbb{R}$, the different planes $\pi_{u,\lambda}$ can be considered as a "sweeping" plane in $\mathbb{R}^3$ and its contact with D and with ∂D will give a method to compute $\chi(D)$:

$$\chi(D) = \sum_{x\in A} \operatorname{sign} K(x) + \frac{1}{2}\sum_{y\in B} \operatorname{sign}(k_g(y) - k_g^u(y)).$$

Here, A is given by the points $x \in D$ such that D is tangent to $\pi_{u,\lambda}$ at x for some λ, B is equal to the subset of points $y \in \partial D$ where ∂D is tangent to $\pi_{u,\lambda} \cap S$ for some λ, and k_g^u denotes the geodesic curvature of $\pi_{u,\lambda} \cap S$. We will show that for a generic $u \in S^2$:

1. the sets A, B are finite;
2. if $x \in A$, then $K(x) \neq 0$;
3. if $y \in B$, then $\pi_{u,\lambda} \cap S$ is a regular curve in S and $k_g(y) \neq k_g^u(y)$.

In this way, the above formula has sense for almost any $u \in S^2$ (that is, for any $u \in S^2 \setminus N$, where N is a null set in S^2). Moreover, this generalizes the result of [2], which computes the Euler-Poincaré characteristic of a plane domain $D \subset \mathbb{R}^2$ by looking at contact with lines.

2. Proof of the formula

The classical Poincaré-Hopf Theorem stays that if S is a closed orientable smooth surface and v is a smooth vector field on S with isolated zeros, then

$$\chi(S) = \sum_{v(x)=0} \operatorname{Ind}_x(v),$$

where $\chi(S)$ is the Euler-Poincaré characteristic of S and the index, $\operatorname{Ind}_x(v)$, is just the local degree of v at x. As an immediate consequence, we get that if $f : S \to \mathbb{R}$ is a Morse

---

The first author is partially supported by Generalitat Valenciana Grant GV-97-CB-12-72. The second author is partially supported by DGICYT Grant PB96–0785.

function (that is, a function with non-degenerate critical points), then

$$\chi(S) = \sum_{x \in \Sigma(f)} \mathrm{Ind}_x(f),$$

where $\Sigma(f)$ denotes the set of singular points of f and the index, $\mathrm{Ind}_x(f)$, is given by $+1$ when x is a local extreme of f, or -1 when x is a saddle point.

The Poincaré-Hopf Theorem was generalized for surfaces with boundary by Morse [4] in the following way: suppose that D is a compact orientable smooth surface with boundary (that we can assume embedded in an orientable smooth surface S without boundary) and let v be a smooth vector field on D such that

1. v has isolated zeros on D,
2. v is not zero on ∂D, and
3. v is tangent to ∂D only at a finite number of points.

If $y \in \partial D$ is a point where v is tangent to ∂D, generically we have that the integral curve of v at y is locally contained in $S \smallsetminus D$ or in D and we can assign an index $+1$ or -1 respectively. In the general case, we can extend this by using the Transversality Theorem and define an index, $\mathrm{Ind}_y(v) \in \{-1, 0, +1\}$. Then, the Morse Theorem stays that

$$\chi(D) = \sum_{v(x)=0} \mathrm{Ind}_x(v) + \frac{1}{2} \sum_{v(y) \in T_y \partial D} \mathrm{Ind}_y(v).$$

Now, we can interpret this in terms of critical points of functions. Suppose that $f : D \to \mathbb{R}$ is a Morse function such that f has no critical points on ∂D and the restriction $f|_{\partial D} : \partial D \to \mathbb{R}$ is also a Morse function. For each point $y \in \Sigma(f|_{\partial D})$, we have that the level set $f^{-1}(f(y))$ is locally contained either in $S \smallsetminus D$ or in D. In the first case, we say that y is a point of type "island" and we assign an index $\mathrm{Ind}_y(f) = +1$ and in the second case, we say that it is point of type "bridge" and assign an index $\mathrm{Ind}_y(f) = -1$ (see Figure 1).



a) "Island" b) "Bridge"

Figure 1

Lemma 2.1. *Let D be a compact orientable smooth surface with boundary and let $f : D \to \mathbb{R}$ be a Morse function such that f has no critical points on ∂D and the restriction $f|_{\partial D} : \partial D \to \mathbb{R}$ is also a Morse function. Then,*

$$\chi(D) = \sum_{x \in \Sigma(f)} \mathrm{Ind}_x(f) + \frac{1}{2} \sum_{y \in \Sigma(f|_{\partial D})} \mathrm{Ind}_y(f).$$

An interesting application of this formula is obtained when $D \subset S \subset \mathbb{R}^3$ and we consider the family of height functions. Given $u \in S^2$, we define the height function $h_u : \mathbb{R}^3 \to \mathbb{R}$ by $h_u(x) = \langle x, u \rangle$. When we restrict this function to D (or S), the level sets are the plane curves $\pi_{u,\lambda} \cap D$ (resp. $\pi_{u,\lambda} \cap S$), where $\pi_{u,\lambda}$ is the plane $h_u^{-1}(\lambda)$. In particular, if λ is a regular value of $h_u|_D$ (and hence of $h_u|_S$), we have that $\pi_{u,\lambda} \cap S$ is in fact a smooth curve in S.

We will give a geometrical interpretation of the critical points of $h_u|_D$ and $h_u|_{\partial D}$ and the corresponding indices. Before that, we need to fix some notation. Given $x \in D$, we denote by $N(x)$ and $K(x)$ the normal vector and the Gauss curvature of D at x respectively. Given $y \in \partial D$, we denote by $n(y)$ and $k_g(y)$ the normal vector and the geodesic curvature of ∂D in D at y respectively (we choose the orientation in ∂D so that $n(y)$ points to the interior of D). Finally, we denote by $k_g^u(y)$ the geodesic curvature of the curve $\pi_{u,\lambda} \cap S$ at a regular point y of $h_u|_D$ (in the case that $\pi_{u,\lambda} \cap S$ and ∂D are tangent at y, we choose in $\pi_{u,\lambda} \cap S$ the same orientation).

Theorem 2.2. *Let $D \subset S \subset \mathbb{R}^3$ be a domain with boundary in an orientable smooth surface S in $\mathbb{R}^3$ and let $u \in S^2$.*

1. *Let $x \in D$. It is a critical point of $h_u|_D$ if and only if $\pi_{u,\lambda}$ is tangent to D at x (where $h_u(x) = \lambda$) if and only if $u = \pm N(x)$.*
2. *Let $x \in D$ be a critical point of $h_u|_D$. It is non-degenerate if and only if $K(x) \neq 0$. Moreover, it is a local extreme of $h_u|_D$ when $K(x) > 0$ and a saddle point when $K(x) > 0$.*
3. *Let $y \in \partial D$ be a regular point of $h_u|_D$. It is a critical point of $h_u|_{\partial D}$ if and only if $\pi_{u,\lambda} \cap S$ is tangent to ∂D at y if and only if $\tilde{u} = \pm n(y)$, where $\tilde{u}$ denotes the normalized orthogonal projection of u in T_yD.*
4. *Let $y \in \partial D$ be a regular point of $h_u|_D$ which is also a critical point of $h_u|_{\partial D}$. It is non-degenerate if and only if $k_g(y) \neq k_g^u(y)$. Moreover, it is an island when $k_g(y) > k_g^u(y)$ and a bridge when $k_g(y) < k_g^u(y)$.*

Proof. Parts (1) and (2) are standard results on Generic Geometry and can be found for instance in [1]. We will prove (3) and (4). Suppose that $\alpha(s)$ is a parametrization of ∂D by arc length, so that $y = \alpha(s_0)$ and $n(y) = N(y) \wedge \alpha'(s_0)$. Since $\{N(y), \alpha'(s_0), n(y)\}$ is a orthonormal basis of $\mathbb{R}^3$, we have that

$$u = \langle u, N(y) \rangle N(y) + \langle u, \alpha'(s_0) \rangle \alpha'(s_0) + \langle u, n(y) \rangle n(y).$$

In particular, provided that $u \neq \pm N(y)$, we have that $\tilde{u} = v / \|v\|$, where

$$v = \langle u, \alpha'(s_0) \rangle \alpha'(s_0) + \langle u, n(y) \rangle n(y).$$

Thus, $\tilde{u} = \pm n(y)$ if and only if $\langle u, \alpha'(s_0) \rangle = 0$ if and only if y is a critical point of $h_u|_{\partial D}$.

On the other hand, let $\beta(s)$ be a parametrization of $\pi_{u,\lambda} \cap S$ by arc length, with $\beta(s_0) = y$. In this case, $\langle u, \beta(s) \rangle = \lambda$ for all s and thus $\langle u, \beta'(s_0) \rangle = 0$. This gives that $\tilde{u}$ is also equal to the normal vector to $\beta'(s_0)$ in $T_y\partial D$. In particular, y is a critical point of $h_u|_{\partial D}$ if and only if the two curves are tangent.

Suppose now that y is a critical point of $h_u|_{\partial D}$. We have that

$$\alpha''(s_0) = k_n(y) N(y) + k_g(y) n(y),$$

where $k_n(y)$ is the normal curvature of α. This gives that

$$\langle u, \alpha''(s_0) \rangle = k_n(y) \langle u, N(y) \rangle + k_g(y) \langle u, n(y) \rangle.$$

But we can do the same with $\beta(s)$, getting

$$0 = \langle u, \beta''(s_0)\rangle = k_n(y)\langle u, N(y)\rangle + k_g^u(y)\langle u, n(y)\rangle.$$

This implies that

$$\langle u, \alpha''(s_0)\rangle = (k_g(y) - k_g^u(y))\langle u, n(y)\rangle.$$

Since $\langle u, n(y)\rangle \neq 0$, we conclude that y is a non-degenerate critical point of $h_u|_{\partial D}$ if and only if $k_g(y) \neq k_g^u(y)$. The fact that the cases $k_g(y) > k_g^u(y)$ and $k_g(y) < k_g^u(y)$ correspond to an island or a bridge respectively can be deduced from the way we have chosen the orientation on ∂D. □

The following consequence can be seen as a stereological version of the Gauss-Bonnet formula.

Corollary 2.3. *Let $D \subset S \subset \mathbb{R}^3$ be a domain with boundary in an orientable smooth surface S in $\mathbb{R}^3$. For a generic $u \in S^2$ the height function $h_u|_D$ is a Morse function, has no critical points on ∂D and its restriction to ∂D is also a Morse function. In particular, we have that*

$$\chi(D) = \sum_{x\in D/N(x)=\pm u} \operatorname{sign} K(x) + \frac{1}{2}\sum_{y\in\partial D/n(y)=\pm\tilde{u}} \operatorname{sign}(k_g(y) - k_g^u(y)).$$

Proof. The formula for $\chi(D)$ is a direct consequence of Lemma 3.1 and Theorem 3.2. We will see the first part of the statement. Given $u \in S^2$, $h_u|_D$ is a Morse function if and only if u is a regular value of the Gauss map $N : D \to S^2$ (just note that the points $x \in D$ where $K(x) = 0$ are the critical points of the Gauss map). Therefore, the Sard Theorem implies that for a generic $u \in S^2$, $h_u|_D$ is a Morse function. The other two conditions are also an easy consequence of this theorem. □

It is also interesting to look at some particular cases. Suppose for instance, that S is a plane, namely $S = \mathbb{R}^2$. Then we can use height functions $h_u : \mathbb{R}^2 \to \mathbb{R}$ for $u \in S^1$, instead of the height functions of $\mathbb{R}^3$. The level curves in this case are the orthogonal lines to u, which have zero curvature. Moreover, h_u has no critical points in $\mathbb{R}^2$ and the geodesic curvature is equal to the classical curvature of a plane curve. Thus, we get the following formula method to obtain the Euler number of a domain in $\mathbb{R}^2$ based on 1-dimensional observations, modified from [3], and that has been used in Stereology, [2].

Corollary 2.4. *Let $D \subset \mathbb{R}^2$ be a domain with boundary. For a generic $u \in S^1$ the restriction of the height function $h_u|_{\partial D}$ is a Morse function and*

$$\chi(D) = \frac{1}{2}\sum_{y\in\partial D/n(y)=\pm u} \operatorname{sign} k(y).$$

In the case that $S = S^2$, the level curves associated to $u \in S^2$ are the corresponding parallels. Note that $h_u|_{S^2}$ is a Morse function with two critical points, u and $-u$. Moreover, if we denote by θ_u, γ_u the spherical coordinates associated to u, we have that the geodesic curvature of the parallel $\gamma_u = \gamma_0$ is $-\tan\gamma_0$.

Corollary 2.5. *Let $D \subset S^2$ be a domain with boundary. For a generic $u \in S^2$,*

$$\chi(D) = \frac{1}{2}\sum_{y\in\partial D/n(y)=\pm\tilde{u}} \operatorname{sign}(k_g(y) + \tan\gamma_u(y)) + \#(\{u, -u\} \cap D),$$

where $\#(\{u, -u\} \cap D)$ means the number of times that u or $-u$ belong to D.

It is also possible to compute the Euler number of $D \subset S^2$ from observations on a meridian that sweeps through D. This is not a consequence of the construction associated to the height functions, but a different family of functions. Let $u \in S^2$ and let S^1 be the circle in S^2 orthogonal to u. Then, we have the regular function $\theta_u : S^2 \smallsetminus \{u, -u\} \to S^1$, whose level sets are the meridians $\theta_u = \theta_0$ (which have zero geodesic curvature). Moreover, for a generic $u \in S^2$, the restriction of θ_u to ∂D is a Morse function, [6].

Corollary 2.6. *Let $D \subset S^2$ be a domain with boundary. For a generic $u \in S^2$,*

$$\chi(D) = \frac{1}{2} \sum_{y \in \partial D / n(y) \perp \tilde{u}} \operatorname{sign} k_g(y) + \#(\{u, -u\} \cap D).$$

Proof. Let $S = S^2 \smallsetminus \{u, -u\}$ and let $\tilde{D}$ be the domain with boundary in S given by $D \smallsetminus (B_1 \cup B_2)$, where B_1, B_2 are small geodesic balls centered at $u, -u$ which do not intersect ∂D. It follows from Lemma 3.1 that

$$\chi(\tilde{D}) = \frac{1}{2} \sum_{y \in \partial D / n(y) \perp \tilde{u}} \operatorname{sign} k_g(y).$$

On the other hand, $\chi(D) = \chi(\tilde{D}) + \#(\{u, -u\} \cap D)$, which concludes the proof. ☐

Remark 2.7. The preceding result may be extended to ovaloids in the following way: Let S be an ovaloid (compact and connected surface in $\mathbb{R}^3$ for which the Gaussian curvature $K > 0$) and $\tilde{D}$ a domain in S; then, S is diffeomorphic to a sphere S^2 through its Gauss map. Let $D \subset S^2$ denote the image of $\tilde{D}$ under this diffeomorphism; then, $\chi(D) = \chi(\tilde{D})$ and it is possible to obtain $\chi(\tilde{D})$ from observations in the curves which are image of the meridians in S^2.

3. Stereological applications

The Euler-Poincaré characteristic for domains in $\mathbb{R}^2$ and $\mathbb{R}^3$ has been studied in several stereological applications (see, for instance [2]). The principle used to obtain the Euler number of an n-dimensional domain in $\mathbb{R}^n$ ($n = 2$ or 3), is based on what happens in an $(n-1)$-dimensional plane that sweeps through the domain. For domains in $\mathbb{R}^2$, this principle is based on Corollary 2.4. Now, we will extend this principle to domains in a surface.

Let $D \subset S \subset \mathbb{R}^3$ be a domain with boundary in an orientable smooth surface S in $\mathbb{R}^3$ and let $u \in S^2$. When λ varies on $\mathbb{R}$, the different planes $\pi_{u,\lambda}$ can be considered as a "sweeping" plane in $\mathbb{R}^3$, and Corollary 2.3 can be expressed as:

$$\chi(D) = (I_2 - B_2) + \frac{1}{2}(I_1 - B_1),$$

where I_1, B_1 denote the number of islands and bridges, respectively, observed in the level curves $\pi_{u,\lambda} \cap S$ (see Figure 2) and I_2, B_2, similarly to [2], denote the number of islands and bridges observed in the "sweeping" plane $\pi_{u,\lambda}$, which contribute to the sum $\sum_{x \in \Sigma(h_u)} \operatorname{Ind}_x(h_u)$ (see Figure 2).



Figure 2

References

[1] J.W. Bruce, P.J. Giblin, *Curves and singularities,* Cambridge University Press, 1984.

[2] H. J. G. Gundersen, R. W. Boyce, J.R. Nyengaard, A. Odgaard, *The conneulor: unbiased estimation of connectivity using physical disectors under projection,* Bone **14** (1993), 217–222.

[3] H. Hadwiger, *Vorlesungen, Über Inhalt, Oberfläche und Isoperimetrie,* Springer-Verlag, Berlin, 1957.

[4] M. Morse, *Singular points of vector fields under general boundary conditions,* American Journal of Mathematics **51** (1929), 165–178.

[5] L. A. Santaló, *Integral Geometry and Geometric Probability,* Addison-Wesley Publishing Company Inc., London, 1976.

[6] E. Teufel, *Differential Topology and the Computation of Total Absolute Curvature,* Math. Ann **258** (1982), 471–480

X. Gual-Arnau, Department de Matemàtiques, Campus Riu Sec, Universitat Jaume I, 12071 Castelló SPAIN

E-mail address: `gual@mat.uji.es`

J.J. Nuño-Ballesteros, Departament de Geometria i Topologia, Universitat de València, Campus de Burjassot, 46100 Burjassot SPAIN

E-mail address: `nuno@uv.es`